

TOWARDS AN UNDERSTANDING OF COMPLEX PHENOMENON: APPLYING PARTIAL ORDER THEORY TO MULTI-INDICATOR SYSTEM

University of Neuchâtel

Department of Sociology

Fbg de l'Hôpital 27, 2000 Neuchâtel, Switzerland

October 26-27, 2018



General role of Multi-Indicator System


Indicators in the framework of partial order


Filomena Maggino


Sapienza - University of Rome
Dep. of Statistical Sciences
Councelor of the Italian Prime Minister

Rainer Bruggemann


Leibniz Institute of Freshwater Ecology and Inland Fisheries Germany








A look at newspapers shows the increasing importance of indicators also in the media.








Often these indicators are used to rank objects, like quality of life in cities


- cities ranked with respect to "quality of life"
- regular reports of the EU, where nations are compared with respect to e.g. innovation potential
- UNICEF: child well-being


- ...






















Indicators allow us to understand a complex system

Indicators allow us to control a complex system

systemic view of multi-indicator system


(2) Role of indicators

indicators and complexity


describing

understanding

controlling





multi-indicator system




(2) Role of indicators indicators and complexity

Consequences of complexity in constructing indicators







(2) Role of indicators

indicators and complexity

Consequences of complexity in constructing indicators

This has methodological consequences





not a compress / pointfold / pointform representation


but


a representation preserving the systemic characteristic of the phenomena


defined by elements and their relationships















































(3) How to obtain indicators

Proper and accurate application of the hierarchical design





complex structure



each indicator measures and represents a distinct component in the description of the phenomenon


(3) How to obtain indicators








(3) How to obtain indicators







































































(5) Indicators and ranking

Preconditions for ranking

- A aim for ranking must be defined
- A suitable set of indicators must be found
- The set of indicators must have the following properties
 - 1. Orientability: accordance with the ranking aim
 - Ordinability: any indicator induces at least a weak order







Partial order based on a MIS has a (mathematical) structure, which can deliver additional insights

Mathematical structure of PO allows data structure to be explored:
separateness / isolation / grids / sensitivity / antichains / CAM-graph (role of fine-tuning)
Marco's approaches

(6) MIS and Poset

Partial order based on a MIS has a (mathematical) structure, which can deliver additional insights


Mathematical structure of PO allows data structure to be explored:

separateness / isolation / grids / sensitivity / antichains / CAM-graph (role of fine-tuning)

Marco's approaches

Question: how many indicators are needed to obtain a reliable partial order

(6) MIS and Poset If indicators are discrete and ordinal with range [o, m(j)]: Two variants: posets based on a) $\Pi[o,m(j)]$, Fattore-approach \rightarrow grid b) $\Pi[o,m(j)]$ realized by objects \rightarrow no grid, but graphtheoretical structure




Hypothesis


- Whether or not a MIS is more than the sum of its constituents depends on the representation in the ranking
- Local representations (parallel coordinates)
 (see e.g. Rocco&Ramirez-Marquez [4]), amoeba (spider) diagrams, glyphs,...)
- Global representation by directed graphs (Hasse diagrams)

Global representation of orders by a directed graph

- Let $q_1, q_2, ..., q_m$ be a MIS and let q_j have at least an ordinal scaling level, then:
- A profile "v" is an ordered set of values v_j of q_j (j=1,..,m):
 (v₁,v₂,...,v_m)
- Two profiles labelled by k and k' fulfill a partial order relation if $v_{ik} \le v_{ik}$ for all j.
- By applying the cover relation a directed graph can be obtained, called a Hasse diagram (HD)



Structures of object-related Hasse diagrams



Both, cannot be seen by considering any single indicator but is a property of the whole MIS

Indicators and their information content

- If a ranking is intended, then the set of indicators bears the relevant information
- Amoeba diagrams (and other techniques) yield just what the numerical values of each single indicator tells us.
- The "object-related" Hasse diagram has a structure which tells us more than any single indicator
- The structure in turn is the result of incomparabilities


Information and incomparabilities, U

- As known, if indicators are selected for a ranking, then they must be orientable and this characteristic induces at least a weak order
- There is a trade-off: one and only one indicator induces at least a weak order:
 U = o , perfect decision, because any object is comparable to any other, no structure.

Information and incomparabilities, U

- However, with many indicators a decision is difficult
- When the number of indicators is large enough (U
 → |X|*(|X|-1)/2)
 no structure, no decision (at least within the level of
 the Hasse diagram itself)





natt: (number of indicators merged in A)

The graph U=f(natt) is always weakly monotonous increasing with natt

Discussion

- Indicator set: a value for its own right
- Typical questions: Latent indicators; Synthetic indicators
- Indicator set for Ranking: MIS
- Representing MIS through an object-related Hasse diagram has a surplus
- Not mentioned: the possibility of poset theory to construct latent indicators

